

02

Territory and environment

694-1700

Environment

Pocket Statistics 2017



Schweizerische Eidgenossenschaft
Confédération suisse
Confederazione Svizzera
Confederaziun svizra

Swiss Confederation


Federal Department of Home Affairs FDHA
Federal Statistical Office FSO

Neuchâtel 2017

- Published by:** Federal Statistical Office (FSO)
- Information:** Laurent Zecha, FSO, tel. +41 58 463 67 20, umwelt@bfs.admin.ch
- Editor:** Laurent Zecha, FSO
In collaboration with the Federal Office for the Environment (FOEN) and the Federal Office of Energy (SFOE)
- Series:** Swiss Statistics
- Topic:** 02 Territory and environment
- Original text:** German
- Translation:** FSO language services
- Layout:** DIAM Section, Prepress/Print
- Graphics:** DIAM Section, Prepress/Print
- Front page:** FSO; Concept: Netthoewel & Gaberthüel, Biel;
Photograph: © Annette Shaff – Fotolia.com
- Printed:** in Switzerland
- Copyright:** FSO, Neuchâtel 2017
Reproduction with mention of source authorised
(except for commercial purposes)
- Orders:** Federal Statistical Office, CH-2010 Neuchâtel,
tel. +41 58 463 60 60, fax +41 58 463 60 61, order@bfs.admin.ch
- Price:** free of charge
- FSO number:** 694-1700



Contents




People change the environment by using natural resources and generating waste and other emissions. On the other hand, environmental conditions also have an effect on people and can prompt them to respond to certain circumstances. The present Pocket Statistics highlights such interactions by means of indicators.

1 Use of natural resources

On the one hand, ecosystems produce services that are useful to people. On the other hand, human activities require natural resources such as water, soil, energy or material. The way in which resources are used affects the supply available, particularly in the case of non-renewable resources.

Consumption of drinking water¹



¹ Corresponds to the quantity from the public water supply (own supply from commerce, industry and agriculture are not included).


Source: Swiss Association of Gas and Water Suppliers

© FSO 2017

Around 80% of drinking water comes from groundwater sources, approximately half of which is spring water. In 2015, 933 million cubic metres of drinking water were produced. The consumption of drinking water in Switzerland has decreased by 20% since 1990.

Drinking water efficiency

Index 1990=100




Sources: Swiss Association of Gas and Water Suppliers;
FSO – SNA, ESPOP/STATPOP

© FSO 2017

The consumption of drinking water has become more efficient since 1990 – the needs of a growing economy were able to be covered by the use of less drinking water. The per capita requirement for drinking water has also decreased: in 1990, daily consumption was 472 litres per person, in 2015, this figure was 307 litres. The water that is used abroad to produce imported products is, however, not included in this figure.

Settlement and urban areas




Source: FSO – Land use statistics

© FSO 2017

7.5% of Switzerland is covered by settlement and urban areas. Within 24 years, the settlement and urban areas have grown by 23% or 584 km², mainly at the expense of agricultural areas. This is equivalent to an area increase of nearly 0.75 m² per second.

Efficiency of use of settlement areas

Index 1979/85=100




Source: FSO – Land use statistics, ESPOP/STATPOP

© FSO 2017

Since the period 1979/85, the settlement and urban areas have grown faster than the population. The settlement and urban area requirement per person has therefore increased: according to the latest figures, this requirement is approximately 407 m² per person, which is roughly 20 m² more than 24 years ago.

Energy use¹




¹ Energy use corresponds to the gross energy consumption minus the import/export balance of electricity which, depending on the year, may be positive or negative and usually accounts for a few percent of the gross consumption. In 2015, more electricity was exported than imported: the energy use was approximately 0.3% higher than the gross consumption. Neither energy use nor gross consumption take into account conversion and distribution losses which can vary greatly depending on the energy carrier. It is assumed, for example, that when producing electricity from hydropower, no losses occur whereas when converting nuclear combustibles, roughly two thirds of the energy escapes in the form of heat that is, however, partly used in district heating.

Source: FSOE – Overall energy statistics

© FSO 2017

In the short term, energy consumption is dictated by the economic situation and weather conditions. In the long term, however, population and economic trends as well as technological and lifestyle changes are also important influencing factors. In 2015, 75% of the energy used in Switzerland came from abroad. Since 1990, gross energy consumption has risen by 6% and in 2015 stood at approximately 1.08 million terajoules.

Electricity consumption




Source: FSOE – Electricity statistics

© FSO 2017

Between 1990 and 2015, electricity consumption in Switzerland increased by 25%, although since the mid-2000s this has been seen to flatten out. In 2015, 60% of electricity was produced by hydropower and 34% by nuclear energy. The remainder came from conventional thermal power plants (4%) and various renewable sources (2.6%), such as biogas, as well as solar and wind energy.

Renewable energies

Share of renewable energies in gross energy consumption




Source: FSOE – Renewable energy statistics

© FSO 2017

In 2015, 20.9% of gross energy consumption came from renewable sources. 63% of renewable energies were obtained from hydropower, followed by wood with 18% and energy generation from renewable waste with 11%. Ambient heat (6%), solar energy (3%), biogas (2%), biofuel (0.9%) and wind energy (0.2%) were involved to a lesser extent in the production of energy. Between 1990 and 2015 the consumption of renewable energies rose by 51%, a greater increase than that seen for total energy consumption which rose by 6% over the same period.

Energy efficiency

Index 1990 = 100



Sources: FSOE – Overall energy statistics; FSO – SNA, ESPOP/STATPOP


© FSO 2017

Since 1990, both the gross domestic product (GDP) and the resident population have grown more considerably than gross energy consumption: In 2015, less energy had to be used to generate one Swiss franc and less energy was used per capita than in 1990. However, this does not include so-called “grey energy”, i. e. the energy that has been used abroad for the production and transport of imported products.

Material footprint

Raw material consumption (RMC)¹

Million tonnes



¹ Estimation


Source: FSO – Environmental accounting

© FSO 2017

Since 2000 Switzerland's material footprint – measured by domestic raw material consumption (RMC) – has risen by approximately 7%. This corresponds to the amount of raw materials extracted in Switzerland or abroad to meet Swiss final demand. In 2014, the material footprint was 140 million tonnes. Over this period raw material extracted in Switzerland accounted on average for 45% of the footprint. Expressed in tonnes of raw material, imports increased by 19% between 2000 and 2014.

Material efficiency

Index 2000 = 100




Source: FSO – Environmental accounting, SNA, ESPOP/STATPOP

© FSO 2017

Switzerland's raw material consumption (RMC) or material footprint was around 17 tonnes per capita in 2014 and was therefore lower than the level measured in 2000. However, the gross domestic product (GDP) has increased more than material consumption since 2000: This means that in 2014, a higher value added was attained per material amount used than in 2000.

Net carbon sink effect of forests

Million tonnes of CO₂



2000: reduced sink effect as result of windstorm Lothar at the end of 1999

Source: FOEN


© FSO 2017

During growth, plants and trees absorb carbon dioxide (CO₂) and ingest carbon (C) in the process. Conversely, when wood is burnt or when trees die or rot, carbon is re-released in the form of CO₂. As in recent years, more wood has grown in Switzerland than is used or dies off, the forest acts as a sink for CO₂.

Pollination

Farmland for which pollination by animals contributes to production

Thousand hectares




Source: FSO

© FSO 2017

In 2015, around 41 000 hectares of cultivated area benefited from pollination by animals, corresponding to 4% of utilised agricultural area and 14% of arable and perennial crop areas. Particularly in the case of fruit or berries, but also rapeseed and sunflowers, production is stimulated by pollinators. The remaining arable land is used mainly for cereal crops whose pollination takes place by the wind.

Leisure time spent outdoors

Share of population



Source: FSO – Omnibus surveys 2011 & 2015

© FSO 2017


In 2015, 45% of the Swiss population indicated that they spent their leisure time outdoors, every day or almost every day. Another 38% do this at least once a week. Only 4% of the population says they spend their leisure time outdoors less than once per month.

2 Emissions and waste

Human activities generate waste and other emissions that enter the air, soil or watercourses. The higher the emissions, the greater the impact on the environment.

Emissions of air pollutants

Thousand tonnes



¹ NO_x comprises NO and NO₂. Emission values are given in NO₂.


Source: FOEN

© FSO 2017

Nitrogen oxides (NO_x) are released during the combustion of fuels, with motorised transport being the main source. NO_x and volatile organic compounds (NMVOC) are precursors for the formation of ozone (O₃) and particulate matter. NO_x together with ammonia (NH₃) lead to the acidification and the over-fertilisation of natural ecosystems such as fens and forests. Emissions of most air pollutants in Switzerland have decreased since the 1990s.

Nitrogen oxide efficiency

Index 1990 = 100




Sources: FOEN; FSO – SNA, ESPOP/STATPOP

© FSO 2017

Since 1990, the gross domestic product (GDP) has grown more considerably than nitrogen oxide (NO_x) emissions, so that nitrogen oxide efficiency has increased. This means that in 2015, less NO_x was emitted per Swiss franc generated than in previous years. The resident population also increased more considerably than nitrogen oxide emissions between 1990 and 2015. Subsequently, less NO_x is released per capita today than several years ago.

Greenhouse gas emissions¹

Million tonnes CO₂ equivalents



¹ According to the FSO air emission accounts which are largely based on the FOEN greenhouse gas inventory. The former are consistent with the National Accounts and for this reason discrepancies occur with respect to the latter.

² HFC, PFC, SF₆, NF₃


Source: FSO – Environmental accounting

© FSO 2017

Due to greenhouse gas emissions, humans intensify the natural greenhouse effect and influence the climate in this way. The majority of these greenhouse gas emissions come from the burning of fossil energy carriers. In 2015, Switzerland emitted around 53 million CO₂-equivalent tonnes (incl. international aviation), 82% of which were due to CO₂ emissions.

Greenhouse gas efficiency

Index 1990 = 100




Source: FSO – Environmental accounting, SNA, ESPOP/STATPOP

© FSO 2017

Since 1990, the gross domestic product (GDP) grew more considerably than greenhouse gas emissions, meaning that greenhouse gas efficiency has increased. The resident population has also increased at a faster pace than greenhouse gas emissions: in 2015, 6.4 CO₂-equivalent tonnes were emitted per person – 2 tonnes fewer than in 1990. However, this does not include so-called “grey emissions”, i. e. emissions created abroad during production and transport of imported products.

Nitrogen balance of agricultural areas

Amounts of nitrogen entering agricultural soils or extracted from them




Source: FSO – Environmental accounting

© FSO 2017

Nitrogen (N) is used as a fertiliser in agriculture. From the excess amount, a part is released into the air as ammonia (NH_3) and another part is leached as nitrate (NO_3) into groundwater. In the last ten years, the nitrogen surplus has been around 90 000 tonnes per year on average.

Phosphorus balance of agricultural areas

Amounts of phosphorus entering agricultural soils or extracted from them




Source: FSO – Environmental accounting

© FSO 2017

Phosphorus (P) is one of the main nutrients of plants and is used in agriculture as a fertiliser. Excess phosphorus can enter surface water through soils and stimulate algae and plant growth, particularly in lakes. When this biomass dies off and decomposes, there may be a shortage of oxygen and thus a disruption of the ecological balance. In the last ten years, excess phosphorus has been around 4 000 tonnes per year on average.

Municipal waste

Million tonnes



■ Separately collected municipal waste¹

■ Municipal waste, incinerated and deposited in landfills²

¹ Total comprising compost, paper and cardboard, glass, tin, aluminium, PET, textiles, batteries (since 1993), electrical and electronic devices (since 2001)

² The figures from 2004 include only domestic waste without imports.


Source: FOEN

© FSO 2017

In 2015, roughly 6 million tonnes of municipal waste was generated in Switzerland. Around 53% of this was separately collected and recycled. In 1990, the corresponding share was 29%. The remainder was burnt in waste incineration plants or (before 2005) burnt or land-filled. The heat generated by incineration is used for district heating or the production of electricity.

Efficiency in municipal waste

Index 1990=100




Sources: FOEN; FSO – SNA, ESPOP/STATPOP

© FSO 2017

The increase in municipal waste since 1990 is roughly in line with the gross domestic product GDP: in 2015, roughly the same amount of waste per Swiss franc generated was caused as in 1990. However, municipal waste has increased more considerably than the resident population: in 2015, municipal waste per capita was 724 kg, around 117 kg more than in 1990. This does not, however, include waste that is generated abroad during the manufacture of imported goods.

Hazardous waste

Thousand tonnes



Source: FOEN

© FSO 2017


Hazardous waste requires special technical and organisational measures for disposal due to its composition and characteristics. Of the approximately 2.3 million tonnes of hazardous waste generated in Switzerland in 2015, 81% was recycled, treated, burned or landfilled, 19% was exported. The increase in recent years is mainly due to contaminated matter excavated during the clean-up of brownfield sites.

3 Environmental situation

The environmental situation is influenced by human activities among other things. In addition to the quality of air, water-courses, soils, ecosystems or landscapes, the supply of these resources is also of interest.

Concentration of particulate matter (PM10)

Annual mean in micrograms per cubic metre



Data prior to 1997 was collected by a different method.

Threshold limit (annual mean): 20 micrograms per cubic metre

Source: FOEN – NABEL


© FSO 2017

Pollution from particulate matter (PM10) has declined in recent years although the daily and yearly thresholds are still partly exceeded, in particular in cities and along heavily used roads. Particulate matter comes from combustion processes, mechanically due to abrasion and resuspension or is formed from precursors. Particulate matter can cause diseases of the respiratory tract and of the cardiovascular system and may also lead to an increase in the risk of cancer.

Ozone layer

Total amount of ozone measured in the air column above Arosa
(annual average values)

Dobson units



Source: MeteoSwiss


© FSO 2017

The stratospheric ozone layer is found at an altitude from 20 to 40 km and protects the earth from UVB rays which can lead to skin damage and even skin cancer among humans. Various substances contribute to the depletion of the ozone layer. The implementation of the Montreal Protocol signed in 1987 for the protection of the ozone layer has been able to reduce the use of such substances. The overall concentration of substances which deplete the ozone-layer in the stratosphere has decreased since 1997.

Mean annual temperature

Deviation from the long-term mean (1961 to 1990)

Degrees Celsius




Source: MeteoSwiss

© FSO 2017

The air temperature varies from year to year and is characterised by colder and warmer periods. Since the start of the 1990s, above average annual temperatures have been recorded in Switzerland: 9 out of 10 of the warmest years since the start of the recording of temperatures were registered in the 21st century and 2015 was the warmest year yet.

Phosphorus levels in selected lakes

Yearly average in micrograms per litre




Source: FOEN – NAWA

© FSO 2017

In particular, phosphorus (P) is used as a fertiliser whereby the excess accumulates in the soil. If the latter is subject to leaching, run-off or soil erosion, phosphorus is discharged into surface waters which can stimulate algae and plant growth, particularly in lakes. In the decomposition of this biomass, there may be a lack of oxygen, presenting a threat to fish and other creatures. In Switzerland, high concentrations of phosphorus are found notably in watercourses in catchment areas with intensive animal production.

Pesticide residues in groundwater, 2013

Share of measuring stations with excessively high concentrations, by main land use



¹ Value required by the Water Protection Ordinance (WPO)


Source: FOEN – NAQUA

© FSO 2017

In 2013, active substances contained in pesticides were detected at 2% of groundwater measuring stations in concentrations that exceed the threshold of the Waters Protection Ordinance of 0.1 microgram per litre. And this threshold was exceeded at 21% of measuring stations for the pesticide degradation products. Agriculture and settlement areas in particular show elevated pesticide concentrations.

Heavy metal soil pollution

Percentage of 97 monitoring stations with at least one measurement above benchmark limits for lead, copper, cadmium or zinc




Source: FOEN/FOAG – NABO

© FSO 2017

Pollutants, such as heavy metals and organic compounds which are difficult to break down, accumulate in the soil where they may obstruct important functions of the soil and enter the food chain through plants. During the survey period 2005 to 2009, the benchmark values were exceeded for at least one heavy metal in 20% of the soil samples tested.

Soil sealing

Proportion of sealed areas as a percentage of total area




Source: FSO – Land use statistics

© FSO 2017

Buildings and roads in particular are considered as sealed soils. Due to the soil sealing, the soil loses its natural ecological function as habitat, reservoir and filter as well as the ability to transform and degrade substances. In the space of 24 years, sealed soils have increased by 29% in Switzerland. According to latest figures, 4.7% of Switzerland's surface area is sealed.

Endangered animals and plants (Red lists)

Status 1994 to 2016, depending on the species group



¹ Reading aid: The endangerment status was assessed for 94% of the 87 mammal species. Data are deficient for the remaining species.

Source: FOEN

© FSO 2017

There are around 46000 known species of flora, fauna and fungi in Switzerland (excluding unicellular or oligocellular organisms). Of the 10699 evaluated species, 35% are on the red list, i.e. they are considered endangered, missing or extinct. At least 49 of the endangered animal and plant species in Switzerland are classified as endangered worldwide.

Populations of breeding birds – Swiss Bird Index®

Index 1990=100




Source: Swiss Ornithological Institute Sempach

© FSO 2017

The populations of bird species that regularly breed in Switzerland (177 species, 174 of which can be evaluated) have increased since 1990. However, the population of the 42 endangered species on the red list decreased over the same period.

Ecological quality of the forest

Part of the forest area with low, medium or high biotope value




Source: WSL – NFI

© FSO 2017

Around a third of Switzerland's surface area is currently covered by forest. Since the survey period 1983/85, the forest area has grown by around 10%. During the same period, the ecological quality of forests has improved: in 87% of forest areas, the biotope value today is rated medium to high. A forest's biotope value is used to evaluate forest populations as a habitat for animals and plants. The diversity of tree species and the structural diversity, for example, are taken into consideration.

Landscape fragmentation

Effective mesh density (number of meshes per 1000 square kilometres), taking account of railway lines, roads down to class 4, settlements, etc.




Source: FOEN – Landscape Monitoring Switzerland

© FSO 2017

The landscape fragmentation through, for example, roads or rail tracks prevents animals from moving freely in nature. This freedom of movement, however, is an important prerequisite for reproduction and thus for maintaining the species. Between 1980 and 2007, the degree of landscape fragmentation in Switzerland increased by 35%. In the Central Plain, where fragmentation is particularly evident, the spaces between these "obstacles" measured an average 0.8 km² in 2007, compared with 310 km² on the southern slopes of the Alps.

Assessment of the environmental quality by the resident population of Switzerland

Share of population



Source: FSO – Omnibus surveys 2011 & 2015

© FSO 2017

In 2015, 95% of the population assessed the quality of the environment in their place of residence as very or quite good. Regarding the quality of the environment in Switzerland overall, 92% of the population had this opinion. These assessments are roughly the same as those from 2011. However, the quality of the environment worldwide was assessed less positively than it had been four years previously. Whereas 23% of the population then said it was very good or quite good, in 2015 only 20% said so.


4 Effects on society

Environmental conditions can affect the quality of life and the health of the population and may also cause damage to infrastructure and buildings or generate other economic costs.

Damage caused by natural events

Floods, debris flows, landslides and rockfall¹

CHF million (adjusted for inflation, base year 2016)



¹ Rockfall taken into account since 2002


Source: WSL

© FSO 2017

Humanity has always been confronted with natural hazards. However, due to the extension of settlement areas in regions at risk and rising property values, the potential damage from natural hazards is also increasing. Flooding, mudflows, landslides or rockfall are influenced, among other factors, by the climatic conditions.

External costs of road and rail traffic

CHF billion



Road traffic

- Goods transport
- Public transport
- Private motorised transport

Rail traffic

- Goods transport
- Passenger transport

Territoriality principle: traffic within Swiss borders


Sources: FSO; ARE

© FSO 2017

External costs are not borne by those responsible but by third parties (often by the general public). In transport, these types of costs are usually incurred in the environmental and health sectors. However, they are also found in costs relating to the damage or depreciation of buildings due to traffic. In 2013, the estimated external costs incurred in Switzerland were CHF 6.7 billion due to road traffic and CHF 0.8 billion due to rail traffic.

Persons affected by noise

Number of persons exposed to noise immissions over the legal limit



¹ Status 2010

² Status 2006


Source: FOEN

© FSO 2017

Noise is an annoying sound which in addition to its impact on health also has an economic and social dimension (e. g. depreciation of property, social segregation). The main source of noise pollution is road traffic. In 2010, about every fifth person (21% of the Swiss population) was exposed at their place of residence to traffic noise that exceeded the legal limit set by the Noise Abatement Ordinance.

Perception of environmental conditions in the place of residence

Share of population




Source: FSO – Omnibus surveys 2011 & 2015

© FSO 2017

In 2015, 24% of the population said they were very or rather disturbed by traffic noise at home with the window open, 19% by air pollution surrounding the house and 10% by radiation from high-voltage power lines or mobile phone antennae. 93% of the population were very or rather satisfied with the landscape surrounding their place of residence. However, the percentage of people who were very satisfied with the landscape in the residential environment fell between 2011 and 2015 from 56% to 49%.

Assessment of the danger for humans and the environment

Share of population



Source: FSO – Omnibus surveys 2011 & 2015

© FSO 2017


Nuclear power stations are seen as the greatest danger for people and the environment. In 2015, 40% of the population said that they were very dangerous. In 2011, 47% were of this opinion. However, it should be noted that the nuclear reactor accident in Fukushima took place one and a half months before the 2011 survey began. With 36%, the loss of diversity in animal and plant species was considered to be the second greatest danger for people and the environment, followed by rising global temperatures due to climate change (34%).

5 Reactions from society

Society can respond to modified environmental conditions, for example, by taking protective measures, creating incentives to reduce pollution or changing behaviour.

Environmentally related taxes revenue

CHF billion, at current prices




Source: FSO – Environmental accounting

© FSO 2017

Taxes are considered as environmental if the taxed object demonstrably has negative effects on the environment, such as motor fuels, for example. The purpose for the introduction of the tax is irrelevant. Between 1990 and 2015, revenue from environmental taxes more than doubled. In 2015 they formed 1.7% of gross domestic product (GDP) and 5.9% of total revenue from taxes and social contributions.

Public environmental protection expenditure

CHF billion, at current prices



¹ Species and landscape protection; since 1993 incl. governmental agri-environment subsidies

² Excl. domestic waste incinerators

2008: introduction of revised financial statistics


Source: FSO – Environmental accounting

© FSO 2017

Environmental protection expenditure reflects the financial expenditure for the prevention, reduction and elimination of pollution or other damage to the environment. In application of the polluter-pays principle, these costs are borne by the polluters. This is largely the case in the sectors of waste and wastewater. Public environmental protection expenditure in Switzerland has increased by 81% since 1990 and in 2014 stood at CHF 4.3 billion.

Protected areas of national importance

Share of national territory (areas with multiple protection counted only once)



¹ Water and migratory bird sanctuaries, federal game reserves, landscapes and natural monuments of national importance


² National park, raised bogs and fenlands, alluvial zones, amphibian breeding areas, moorland, dry meadows and pastures

Source: FOEN

© FSO 2017

Areas of national importance for the protection of biodiversity and the landscape together cover roughly 23% of Switzerland's surface area. 4% of the surface area is strictly protected, i.e. its undiminished preserved state is enshrined in law.

Modal split in passenger and goods transport



¹ As percentage of tonne-kilometres


² As percentage of person-kilometres

Source: FSO

© FSO 2017

Both goods and passenger transport performance have increased in recent years. However, the share of goods transport by rail has decreased. In 2015, the corresponding share was 38.6%. In contrast, the share of passenger transport in public transport steadily increased between 1998 and 2011. Since then it has declined slightly and in 2015 amounted to approximately 20.7%.

Average CO₂ emissions from new passenger cars



Values before 2002 were calculated using another method.

According to the CO₂-Act, average emissions should be reduced to 130 grammes per kilometer by 2015.


Source: SFOE

© FSO 2017

According to test-bench measurements, average CO₂ emissions from new passenger cars fell between 1996 and 2016 by 38% and in 2016 were 134 g per kilometre. However, the CO₂ emissions from all passenger cars dropped by only 3.9% between 1996 and 2015. This difference is due to the inclusion of older passenger cars, the increase in traffic volume as well as the growing gap between test-bench measured CO₂ emissions and those actually arising in road traffic. On average this gap was 13% in 2005. By 2015 it had risen to 41%.

Separately collected municipal waste (recycling)

Collection rate



¹ A deposit can be introduced if the recycling rate of 75% stipulated in the Beverage Container Ordinance (BCO) is not achieved.


Source: FOEN

© FSO 2017

In 2015, 382 kg of waste from households and industry per person was collected separately and entered the recycling chain. This corresponds to approximately 53% of the total municipal waste. Scrap paper, green waste and glass make up the largest quantities here.

Consumption of organic products

Share of private household expenditure on products with an organic label out of the total expenditure for food and drink




Source: FSO – HBS

© FSO 2017

Products that are produced in accordance with the ordinance on organic farming are labelled as organic. Among other things, the principles apply that no chemical-synthetic pesticides and no modified organisms are used and that livestock are kept according to specific regulations. In 2014, around 8% of the total expenditure on food and drink was spent on organic products.

Environmental behaviour in everyday life

Share of population




Source: FSO – Omnibus surveys 2011 & 2015

© FSO 2017

In 2015 the population was more likely than four years previously to take electricity consumption into account when purchasing small electrical devices or lamps. In 2015, 42% said they always consider this, compared with 35% in 2011. In 2015 when asked if they reduce the heating when their dwelling is empty for at least two days, 44% said they “always” did so and 20% “never”. When asked about their consumption of organically produced food, another environmentally relevant aspect of everyday life, 7% of the population said they bought only organic products and the same percentage never do so.

Environment-related patents

Share of environment-related patents of the total number of patents filed by Swiss inventors



Source: OECD

© FSO 2017

In 2013, 237 environment-related patents were filed by Swiss inventors. As a percentage of all filed patents, this corresponds to a share of approximately 7.4%. Environment-related patents include innovations in the sectors of renewable energy production, energy efficiency, environmental management and emission-reducing technologies.

Comparison of some European countries

	AUT	BEL	CH	DEU	DNK	ESP
Share of forest areas, 2014	46.9%	22.6%	31.6%	32.7%	14.4%	36.8%
Share of organic farming area in total utilised agricultural area, 2015	20.3%	5.1%	13.0%	6.4%	6.3%	8.2%
Gross energy consumption, 2015 (t oil equivalent/person)	3.8	4.8	3.1	3.8	2.9	2.6
Share of renewable energies in gross energy consumption, 2015	29.0%	6.7%	20.9%	12.2%	28.4%	13.7%
Emissions of greenhouse gases, 2014 (t/person)	8.9	10.1	5.9	11.1	9.0	7.1
Emissions of nitrogen oxide, 2014 (kg/person)	17.6	17.5	8.3	15.1	20.0	17.3
Domestic material consumption (DMC), 2015 (t/person)	21.9	12.9	11.3	16.1	19.8	8.3
Household waste, 2015 (kg/person)	560	418	725	625	789	434
Share of separately collected municipal waste (recycling and compost), 2015	56.9%	53.4%	52.7%	66.1%	46.3%	33.3%
Average CO ₂ emissions from new passenger cars, 2014 (g/km)	129	121	142	133	110	119
Share of environmental taxes in total tax and social contribution revenue, 2015	5.5%	4.7%	5.9%	5.0%	8.6%	5.6%
Threatened bird species as percentage of all known bird species ¹	27%	20%	35%	36%	16%	21%

¹ Latest data available ² Status 2012 ³ Status 2014 ⁴ Threatened indigenous species

Sources: FSO; Eurostat; OECD

FIN	FRA	GBR	GRC	IRL	ITA	LUX	NLD	PRT	SWE	EU-28
73.1%	30.8%	12.9%	31.2%	10.9%	31.4%	33.5%	11.1%	34.9%	68.9%	...
9.9%	4.8%	2.9%	8.4%	1.7%	11.8%	3.2%	2.7%	6.5%	17.1%	5.9% ³
6.0	3.8	2.9	2.3	3.0	2.6	7.2	4.6	2.2	4.6	3.2
31.6%	8.6%	7.7%	11.3%	7.6%	16.8%	4.9%	4.7%	21.6%	42.2%	13.0%
10.8	6.9	8.1	9.3	12.6	6.9	19.1	11.1	6.2	5.6	8.4
25.1	13.3	14.6	22.8	16.6	13.0	50.0	13.9	15.4	13.9	15.4
30.5	11.3	8.8	12.7	20.7	6.9	21.9	11.1	15.8	22.5	13.2
500	501	485	506 ²	587 ²	486	625	523	453 ³	447	476
40.6%	39.5%	43.5%	19.3% ²	36.6% ²	43.5%	48.0%	51.7%	30.4% ³	48.0%	45.0%
127	114	125	108	117	118	130	107	109	131	125
6.7%	4.8%	7.4%	10.3%	8.0%	7.9%	4.9%	9.0%	7.0%	5.1%	6.3%
24%	27%	2% ⁴	14%	24%	28%	19%	22%	28%	20%	...

Glossary

Biodiversity Biodiversity spans the various species of animals, plants, fungi and bacteria, genetic diversity within the species and the different habitats in which the species live.

CO₂-equivalents Emissions of greenhouse gas emissions other than CO₂ (CH₄, N₂O, HFKW, PFKW and SF₆ and NF₃) are converted into CO₂-equivalents according to their global warming potential (GWP) to ensure better comparability. 1 kg CH₄ corresponds to 25 kg CO₂, 1 kg N₂O corresponds to 298 kg CO₂.

Eco-efficiency Measure for the economic performance (e. g. GDP) in relation to the environmental impacts (e. g. the resources used or the emissions released).

Emissions The release of pollutants, noise or radiation from natural sources or by humans into the environment.

Energy carriers Any substance which can be used to generate energy, whether directly or after transformation. Fossil energy carriers are all primary energy carriers which have come from organic substances in the soil (petroleum, natural gas, various hydrocarbons, coal etc.)

Energy use Corresponds to the gross energy consumption minus the electricity import/export balance.

External costs Costs incurred during production or consumption that are not borne by those responsible but a third party.

GDP (Gross Domestic Product) GDP is used for measuring a country's economic performance over a period of one year.

Greenhouse effect The greenhouse effect is a natural phenomenon. It originates from various gases in the atmosphere (water vapour, carbon dioxide, methane, nitrous oxide etc.) which reflect part of the heat radiation leaving the earth. An increase in the concentration of such greenhouse gases leads to the heating of the atmosphere.

Greenhouse gases Natural or man-made gaseous substances in the air which contribute to the greenhouse effect. The Kyoto Protocol considers the following greenhouse gases or groups of gases: carbon dioxide (CO₂), methane (CH₄), nitrous oxide (N₂O), hydrofluorocarbons (HFCs), perfluorocarbons (PFCs), sulphur hexafluoride (SF₆) and nitrogen trifluoride (NF₃).

Gross energy consumption The gross energy consumption consists of the domestically produced primary energy, the foreign trade totals of various energy carriers as well as changes in stocks.

Immissions Levels of air pollutants, noise, vibration or radiation at their point of impact.

Limit values Limit values are used to evaluate harmful or noxious impacts. They consider the effects of emissions on the environment and more sensitive groups of people such as children, the sick, the elderly and pregnant women. Thresholds are defined in terms of air pollution, noise pollution, vibration and radiation.

NMVO (**N**on **M**ethane **V**olatile **O**rganic **C**ompounds) Volatile organic compounds that exclude methane and FCKW. They are precursors for the formation of ozone, summer smog and PM10.

PM10 (**P**articulate **M**atter <10 μm) Dust particles with a diameter of less than 10 microns. These particles are caused by combustion processes, mechanically due to abrasion and resuspension or are formed from precursors. The main sources of PM10 are motorised transport, agriculture and forestry, industry and commerce (including construction sites).

Renewable energy Energy sources that are available for an unlimited period in principle. This includes the use of hydropower, solar energy, ambient heat, biomass, and wind energy, renewable energy from waste as well as energy from waste water treatment plants.

Further information

Federal Statistical Office (FSO) website:

www.statistics.admin.ch → Look for statistics → Territory and environment

Data relating to the graphs shown:

www.statistik.ch → Statistiken finden → Raum, Umwelt → Umwelt und Ressourcen

NewsMail subscription:

www.statistik.ch → Dienstleistungen → NewsMail-Abonnement

Federal Office for the Environment (FOEN): www.foen.admin.ch

Swiss Federal Office of Energy (SFOE): www.sfoe.admin.ch

MeteoSwiss: www.meteoswiss.ch

Orders

Tel. 058 463 60 60

Fax 058 463 60 61

order@bfs.admin.ch

Price

Free of charge

FSO number

694-1700

Statistics
counts for you.

www.statistics-counts.ch